



Entrez	PubMed	Nucleotide	Protein	Genome	Structure	PMC	Journals	Books
Search	PubMed	▼ for					Go	Clear
<input checked="" type="checkbox"/> Limits Preview/Index History Clipboard Details								

Display	Abstract	▼	Show: 20	▼	Sort	▼	Send to	Text	▼
---------	----------	---	----------	---	------	---	---------	------	---

☐ 1: Indian J Biochem Biophys. 1993 Dec;30(6):389-94.

[Related Articles, Links](#)

Entrez  
PubMed

## Is asparagine-linked protein glycosylation an obligatory requirement for angiogenesis?

**Banerjee DK, Vendrell-Ramos M.**

PubMed  
Services

Department of Biochemistry, School of Medicine, University of Puerto Rico, San Juan 00936-5067.

Related  
Resources

Dependence of protein N-glycosylation on capillary endothelial cell proliferation has been studied. Amphomycin, a potent N-glycosylation inhibitor, inhibited capillary endothelial cell proliferation in a dose-dependent manner. beta-Agonist isoproterenol as well as other intracellular cAMP enhancing agents, viz. cholera toxin, prostaglandin E1 and 8Br-cAMP, also enhanced capillary endothelial cell proliferation. In addition to cell proliferation, isoproterenol also enhanced protein glycosylation in these cells. Isoproterenol effect was mediated by beta-adrenoreceptors, as it got reduced on pre-treatment of cells with either atenolol or ICI 118, 551 or propranolol. Furthermore, isoproterenol stimulation of protein glycosylation by exogenous dolichyl monophosphate and its inhibition by tunicamycin (GlcNAc-1P transferase inhibitor) supported the concept that isoproterenol specifically stimulated protein N-glycosylation event(s) in the cell.

PMID: 7516308 [PubMed - indexed for MEDLINE]

Display	Abstract	▼	Show: 20	▼	Sort	▼	Send to	Text	▼
---------	----------	---	----------	---	------	---	---------	------	---

[Write to the Help Desk](#)  
[NCBI | NLM | NIH](#)  
[Department of Health & Human Services](#)  
[Freedom of Information Act](#) | [Disclaimer](#)